

**RECENT PROGRESS IN CHALCONE DERIVATIVES: SYNTHETIC STRATEGIES,
PHARMACOLOGICAL ACTIVITIES AND MOLECULAR INSIGHTS****Neha Pisal***

Hinjewadi Phase 1, Pune Maharashtra, India.

***Corresponding Author: Neha Pisal**

Hinjewadi Phase 1, Pune Maharashtra, India.

DOI: <https://doi.org/10.5281/zenodo.20444912>**How to cite this Article:** Neha Pisal* (2026). Recent Progress In Chalcone Derivatives: Synthetic Strategies, Pharmacological Activities And Molecular Insights. World Journal of Pharmaceutical and Medical Research, 12(6), 71–79.
This work is licensed under Creative Commons Attribution 4.0 International license.

Article Received on 28/04/2026

Article Revised on 18/05/2026

Article Published on 01/06/2026

ABSTRACT

Chalcones represent an important class of flavonoid-related compounds that occur in both natural and synthetic forms. They are structurally characterized by an α , β - unsaturated carbonyl framework, which plays a crucial role in their chemical reactivity and pharmacological potential.^[6,7] These compounds are widely distributed in plants and are involved in various physiological processes, including defense mechanisms and metabolic functions.^[13] Owing to their simple structure and ease of modification, chalcones have gained significant attention as promising scaffolds in medicinal chemistry and drug discovery.^[17,21] Conventional synthetic approaches such as Claisen-Schmidt condensation remain widely used; however, recent advancements have introduced modern techniques including micro- assisted synthesis and green chemistry approaches that improve reaction efficiency, reduce time, and minimize environmental impact.^[1,2,14] Chalcone derivatives exhibit a wide range of biological activities, including antimicrobial, anti-inflammatory, antioxidant, anticancer, and antitubercular effects. These properties are primarily attributed to the presence of an electrophilic α , β - unsaturated carbonyl group, which facilitates interaction with biomolecular targets such as enzymes and proteins.^[6,12] Furthermore, computational approaches such as molecular docking and in silico studies have enhanced the understanding of ligand-target interactions, thereby supporting rational drug design.^[60-62] This review provides a comprehensive overview of recent developments in chalcone chemistry, including synthesis, characterization, biological activities, and mechanisms of action. Special emphasis is placed on structure-activity relationship (SAR) and the influence of substituents on biological properties. The future potential of chalcone derivatives in pharmaceutical research is also highlighted.

KEYWORDS: Chalcone; Claisen –Schmidt condensation; Microwave synthesis; Green chemistry; Antimicrobial; Molecular docking; SAR.**1. INTRODUCTION**

Chalcone are an important class of bioactive compounds belonging to the flavonoid family, chemically defined as 1,3-diary-2-propen-1-one derivatives.^[6,7] their structure consists of two aromatic rings linked by a three-carbon α , β -unsaturated carbonyl system, which is responsible for their unique physicochemical and biological properties. Due to their structural simplicity and high reactivity, chalcones serve as valuable intermediates in organic synthesis and are widely explored as pharmacologically active scaffolds.^[17,21]

Naturally occurring chalcones are found in a variety of plant sources such as liquorice and hops, where they act as key intermediates in the biosynthesis of flavonoids and related compounds.^[13,17] These molecules contribute

to plant defense mechanisms by exhibiting antimicrobial and antioxidant activities. Their diverse pharmacological potential has attracted considerable attention in recent years, particularly due to their ability to interact with multiple biological targets.^[13]


Numerous studies have reported that chalcone derivatives possess a broad spectrum of biological activities, including antimicrobial, anti-inflammatory, anticancer, and antiviral properties.^[6,16] These activities are largely associated with the α , β -unsaturated carbonyl moiety, which acts as an electrophilic centre capable of reacting with nucleophilic sites in proteins and enzymes, thereby modulating various cellular pathways.^[12]

Recent developments in chalcone research have focused on improving synthetic strategies and expanding their pharmacological applications. Traditional methods are increasingly being complemented by modern approaches such as microwave-assisted synthesis and environmentally friendly techniques, which offer advantages such as reduced reaction time, improved yields, and sustainability.^[1,2] In addition, computational tools such as molecular docking and quantitative structure – activity relationship (QSAR) studies have provided deeper insights into chalcone-target interactions, facilitating the design of more potent derivatives.^[30,60]

This review aims to present a detailed overview of chalcone chemistry, including synthesis, characterization, biological activities, mechanism of action, and structure-activity relationships, with a focus on their potential applications in drug discovery.

2. CHEMISTRY OF CHALCONES

Chalcones are characterized by the presence of an α , β -unsaturated carbonyl system, which plays a crucial role in their reactivity and biological activity. The general structure can be represented as.



where Ar and Ar' are aromatic rings.^[7]

2.1 Structural Features

The conjugated system present in chalcones allows electron delocalization the molecule, enhancing stability

and reactivity. This conjugation is responsible for their UV-visible absorption and characteristic coloration was explain.^[7]

2.2 Functional Groups and Reactivity

- **Carbonyl group (C=O):** participates in nucleophilic addition
- **Double bond (C=C):** acts as electrophilic centre
- **Aromatic rings:** allow substitution and structural modification

The α , β -unsaturated system enables chalcones to act as Michael acceptors, which essential for their biological activity.^[6,12]

2.3 Chemical Reactions

Chalcones undergo several reactions

- Michael addition
- Cyclization to flavonoids
- Oxidation and reduction
- Substitution reactions

These reactions make chalcones versatile intermediates in organic synthesis.^[45]

2.4 Stereochemistry

Chalcones predominantly exist in the **trans (E)-form**, which is more stable and biologically active due to reduced hindrance.^[27]




Fig.1: General structure of chalcone showing α , β -unsaturated carbonyl system.

3. METHODS OF SYNTHESIS

3.1 Claisen–Schmidt Condensation

The Claisen–Schmidt condensation is the most widely used method for chalcones synthesis.^[45,46]

Reaction

Aromatic aldehyde + Aromatic ketone → Chalcone.
(Fig.no. 2)

This method is simple and provides high yields.

Mechanism

1. Formation of enolate ion
2. Nucleophilic attack aldehyde
3. Formation of β -hydroxy ketone
4. Dehydration




Fig. 2: Claisen – Schmitt condensation mechanism for chalcone synthesis.

3.2 Microwave-Assisted synthesis

Microwave irradiation significantly reduces reaction time and improves yield.^[1,9] It enhances molecular collisions and reaction efficiency.

Studies have shown.

- Faster reactions (minutes vs hours)
- Higher purity products
- Reduced solvent use.^[3,32,34]

3.3 Green Synthesis

Green chemistry approaches focus on eco-friendly synthesis using.

- Non-toxic solvents

- Reusable catalysts
- Energy-efficient methods.^[14,33]

MicellRESUar media and ionic liquids have been successfully used for chalcones synthesis.^[14]

3.4 Solvent-Free Synthesis

Solvent-free methods reduce environmental impact and improve reaction efficiency. These methods are widely used in microwave-assisted synthesis.^[35]




Fig.3: Modern synthetic approaches for chalcone derivatives.

4. CHARACTERIZATION OF CHALCONES

Characterization of chalcone derivatives is essential to confirm their chemical structure, purity, and functional group composition. Various spectroscopic techniques such as IR, NMR, and Mass spectroscopy are commonly employed to establish the identity of synthesized chalcones.^[51,54]

4.1 Infrared (IR) Spectroscopy

IR spectroscopy is a powerful technique used to identify functional groups present in chalcone based on their characteristic vibrational frequencies.

IR confirms

- Presence of carbonyl group

- Conjugated system
- Functional group substitutions

Key Functional Groups peaks in Chalcones

1. Carbonyl Groups (C=O Stretching)

Range :1650-1680 cm^{-1}

- This is the most important peak for chalcone.
- Appears slightly lower than normal ketone ($\sim 1715 \text{ cm}^{-1}$) due to conjugation with C=C double bond.
- Confirms presence of α,β - unsaturated ketone system.

Interpretation

Lower frequency = strong conjugation \rightarrow confirms chalcone structure.^[51,52]

2. Olefinic Double Bond (C=C Stretching)

Range: $\sim 1600 \text{ cm}^{-1}$

- Represents α, β -unsaturated double bond.
 - Confirms formation of enone system.
- If peak is strong \rightarrow good conjugation.^[51-54]

3. Aromatic C-H Stretching

Range: $\sim 3000- 3100 \text{ cm}^{-1}$

- Indicates presence of benzene rings
- Usually appears as multiple peaks

4. Additional Peaks

- -OH group: $\sim 3200-3500 \text{ cm}^{-1}$ (broad peak)
- -OCH₃ group: $\sim 2830-2850 \text{ cm}^{-1}$
- -NO₂ group: $1500-1550 \text{ cm}^{-1}$

These peaks help identify substituted chalcone derivatives.

4.2 NMR Spectroscopy

NMR spectroscopy provides detailed structural information about chalcones, including proton environment and carbon skeleton.

- Confirms double bond formation
- Confirms trans configuration
- Identifies substituents.^[52]

4.2.1 ¹H NMR (Proton NMR)

Typical Range: $\delta 6-8 \text{ ppm}$

1. Olefinic Protons (-CH=CH-)

- Appear at $\delta 6.5 - 7.5 \text{ ppm}$
- Show doublet signals due to coupling
- Large coupling constant ($J \approx 15-16 \text{ Hz}$). Confirms trans (E)- configuration.

2. Aromatic Protons

- Appear at $\delta 7 - 8 \text{ ppm}$
- Multiplet pattern
- Confirms presence of benzene rings

3. Substituent Peaks

- -OH proton: $\delta 9-12 \text{ ppm}$ (broad)
- -OCH₃ proton: $\delta \sim 3.5-4 \text{ ppm}$

4.2.2 ¹³C NMR (Carbon NMR)

- Confirms carbon framework
- Identifies functional groups
- Supports IR findings

Key Range: $\delta 190-200 \text{ ppm}$

1. Carbonyl Carbon (C=O)

- Appears at $\delta 120-150 \text{ ppm}$

2. Olefinic Carbons

- Appear at $\delta 120-150 \text{ ppm}$

3. Aromatic Carbons

Appear at $\delta 110-140 \text{ ppm}$

4. Substituent Carbons

- -OCH₃: $\delta \sim 55-60 \text{ ppm}$

4.3 Mass Spectrometry

Mass spectrometry is used to determine.

- Molecular weight
- Molecular formula
- Fragmentation pattern

1. Molecular Ion Peak (M⁺)

- Represents exact molecular weight of chalcone
- Confirms successful synthesis

2. Fragmentation Pattern

Chalcones show characteristic fragmentation.

Common Fragmentations

- Loss of CO group
- Cleavage at α,β - double bond
- Formation of benzoyl fragments

Example Fragmentation Flow

Chalcone \rightarrow Break at C=C \rightarrow

Aromatic fragment + carbonyl fragment

3. Base Peak

- Most intense peak
- Represents most stable fragment.

Importance of MS

- Confirms molecular structure
- Detect impurities
- Supports IR and NMR data.^[51-54]

o Overall summary of characterization.

Sr.no	Technique	Information Obtained
1	IR	Functional groups
2	¹ HNMR	Proton environment
3	¹³ CNMR	Carbon structure
4	MS	Molecular weight & fragmentation

5. BIOLOGICAL ACTIVITIES OF CHALCONES

Chalcones exhibit **broad-spectrum pharmacological activities** due to their reactive α , β unsaturated carbonyl system and structural flexibility.^[6,16,21] Their biological effects arise from interaction with enzymes, proteins, and cellular pathways.

5.1 Antimicrobial Activity

Chalcones exhibit antimicrobial activity by.

- Disrupting cell membranes
- Inhibiting enzymes
- Interfering DNA replication

Explanation

- Chalcones increase membrane permeability, leading to leakage of ions and proteins.
- Inhibit bacterial enzymes such as DNA gyrase → prevents replication.
- Interfere with protein synthesis.

Studies confirm activity against Gram-positive and Gram-negative bacteria.^[25,41,66]

5.2 Anti-Inflammatory Activity

Chalcones inhibit inflammatory mediators such as COX and LOX enzymes, reducing prostaglandin synthesis.

Explanation

- Chalcones inhibit cyclooxygenase (COX) and lipoxygenase (LOX) pathway.
- Reduce cytokine production
- Block TNF – α signalling pathway.^[47]

5.3 Antioxidant Activity

Chalcones act as a free radical scavenger, neutralizing ROS and preventing stress.

Explanation

- Phenolic groups donate electrons /hydrogen
- Stabilize free radicals
- Prevent lipid peroxidation.^[20]

5.4 Anticancer Activity

Chalcones are multi-target anticancer agents.

Mechanisms include.

- Induction of apoptosis
- Cell cycle arrest
- Inhibition of angiogenesis
- Proteasome inhibition by chalcones in cancer cells.^[12]

Explanation

- Induce apoptosis via mitochondrial pathway
- Arrest cell cycle at G2/M phase
- Inhibit angiogenesis and tumour proliferation.^[23]

5.5 Antitubercular Activity

Chalcones inhibit Mycobacterium tuberculosis by interfering with essential enzymes.

Chalcone → inhibits mycobacterial enzymes → disrupts cell wall synthesis → inhibits bacterial growth.^[27]




Fig.4: Biological activities and mechanisms of action.

6. MECHANISM OF ACTION

The biological activity of chalcones is primarily due to the α , β -unsaturated carbonyl group, which acts as a Michael acceptor.^[12]

Mechanisms include.

- Enzyme inhibition
- Protein interaction

- DNA binding
- ROS modulation

7. STRUCTURE-ACTIVITY RELATIONSHIP (SAR)

SAR studies show that biological activity depends on substituents.

7.1 Effect of substituents

i. Electron Donating Groups (EDG)

Examples: $-\text{OH}$, $-\text{OCH}_3$

- Increase antioxidant activity
- Enhance hydrogen bonding
- Improve radical scavenging

ii. Electron Withdrawing Groups (EWG)

Examples: $-\text{NO}_2$, $-\text{Cl}$, $-\text{F}$

- Increase antimicrobial and anticancer activity
- Enhance electrophilicity.^[30]

7.2 Position of substitution

Position	Effect
Para	Highest activity
Meta	Moderate
Ortho	Steric hindrance

7.3 Conjugation Effect

- More conjugation \rightarrow higher activity
- Stabilizes molecule
- Enhances binding with proteins

7.4 Lipophilicity

- Increased lipophilicity \rightarrow better membrane penetration
- Improves drug absorption

7.5 Heterocyclic substitution

- Enhances biological activity
- Improves selectivity.^[18]




Fig. 5: SAR of Chalcone.

8. MOLECULAR DOCKING STUDIES

Docking studies is essential for drug design and target identification.

Docking predicts how chalcones bind to proteins.

Importance

- Predicts biological activity
- Save time and cost
- Helps in drug optimization.^[60]

8.1 Software used

- Chem Sketch^[60]
- PyRx^[61]

- SwissADME^[62]

These studies predict binding affinity and pharmacokinetics.

8.2 Docking process

Ligand (chalcone) + Protein (1KZN, 4WMZ)

↓
Binding interaction

↓
Calculation of binding energy

9. APPLICATIONS OF CHALCONES

Chalcones have wide applications.

9.1 Pharmaceutical Applications

- Antimicrobial drugs
- Anticancer agents
- Anti-inflammatory drugs

9.2 Cosmetics

- Antioxidant creams
- Anti-aging formulations

9.3 Agriculture

- Pesticides
- Plant growth regulators

9.4 Chemical Industry

- Intermediate in synthesis
- Dye production.^[15]

10. FUTURE PERSPECTIVES

Future research directions include.

10.1 Drug Development

- New derivatives with better activity
- Target- specific drugs

10.2 Green Chemistry

- Eco-friendly synthesis
- Sustainable production

10.3 Clinical Studies

- Human trials
- Safety evaluation

10.4 Computational Approaches

- AI – based drug design
- Advanced docking studies

11. CONCLUSION

Chalcones represent a highly versatile and promising class of compounds in medicinal chemistry. Their simple structure, ease of synthesis, and diverse biological activities make them valuable scaffolds for drug development. Advances in synthesis techniques, particularly microwave-assisted and green chemistry approaches, have improved efficiency and sustainability. The presence of the α,β -unsaturated carbonyl system plays a crucial role in their biological activity through mechanisms such as enzyme inhibition and protein interaction. Structure– activity relationship studies have further enhanced understanding of how substituents influence activity. Additionally, molecular docking studies have provided insights into target interactions, facilitating rational drug design. Future research focusing on clinical studies, green synthesis, and computational approaches will further expand the therapeutic potential of chalcone derivatives.

12. REFERENCES

1. Gangrade D, Lad S, Mehta A. Overview on microwave synthesis: important tool for green chemistry. *Int J Res Pharm Sci*, 2015; 5(2): 37–42.
2. Polshettiwar V, Varma RS. Microwave-assisted organic synthesis and transformations using benign reaction media. *Acc Chem Res*, 2008; 41: 629–639.
3. Shweta S, Prakash C, Kumud I. Microwave irradiation synthesis of various substituted chalcones using heterogeneous catalysts under solvent-free conditions and their biological studies. *Chem Sci Trans*, 2013; 2: 343–348.
4. Srivastava YK. Eco-friendly microwave-assisted synthesis of some chalcones. *Rasayan J Chem*, 2008; 1(4): 884–886.
5. Jayapal MR, Sreedhar NY. Anhydrous K_2CO_3 as catalyst for the synthesis of chalcones under microwave irradiation. *J Pharm Sci Res*, 2010; 2: 644–647.
6. Dimmock JR, Elias DW, Beazely MA, Kandepu NM. Bioactivity of chalcones. *Curr Med Chem*. 1999; 6: 1125–1149.
7. Dhar DN. *The chemistry of chalcones and related compounds*. New York: Wiley; 1981.
8. Belsare DP, Kazi A. Microwave-assisted synthesis of flavones and their comparative study with conventional method. *IOSR J Pharm*, 2013; 4: 23–27.
9. Caddick S. Microwave-assisted organic reactions. *Tetrahedron*. 1995; 51: 10403– 10432.
10. Kamboj RC, Arora R, Sharma G, Kumar D, Sharma C, Joshi R, Aneja KR. Ecofriendly synthesis and antimicrobial activity of chalcones. *Der Pharma Chem*, 2010; 2(3): 157–170.
11. Shalaby MA, et al. Synthesis, reactions and application of chalcones: a systematic review. *Org Biomol Chem*, 2023.
12. Bazzaro M, Anchoori RK, Mudiam MKR, Issaenko O, Kumar S, Karanam B, Lin Z, Isaksson Vogel R, Gavioli R, Destro F, et al. α,β -Unsaturated carbonyl system of chalcone-based derivatives is responsible for broad inhibition of proteasomal activity and preferential killing of HPV-positive cervical cancer cells. *J Med Chem*, 2010; 54(2): 449–456.
13. Dhaliwal JS, Moshawih S, Goh KW, Loy MJ, Hossain MS, Hermansyah A, Kotra V, Kifli N, Goh HP, Dhaliwal SKS, et al. Pharmacotherapeutic applications and chemistry of chalcone derivatives. *Molecules*, 2022; 27(20): 7062.
14. Dotta D, Gastaldi M, Fin A, Barbero N, Barolo C, Cardano F, Rossi F, Brunelli F, Viscardi G, Tron GC, et al. Chalcone synthesis by green Claisen–Schmidt reaction in cationic and nonionic micellar media. *J Org Chem*, 2025; 90(5): 2915–2926.
15. Dziągwa-Becker M, Oleszek M, Zielińska S, Oleszek W. Chalcones: features, identification techniques, attributes and application in agriculture. *Molecules*, 2024; 29(10): 2247.
16. Elkanzi NAA, Hrichi H, Alolayan RA, Derafa W, Zahou FM, Bakr RB. Synthesis of chalcone

- derivatives and their biological activities: a review. *ACS Omega*, 2022; 7(32): 27769–27786.
17. Gomes MN, Muratov EN, Pereira M, Peixoto JC, Rosseto LP, Cravo PVL, Andrade CH, Neves BJ. Chalcone derivatives: promising starting points for drug design. *Molecules*, 2017; 22(8): 1210.
 18. Mezgebe K, Melaku Y, Mulugeta E. Synthesis and pharmacological activities of chalcone and its derivatives bearing N-heterocyclic scaffolds: a review. *ACS Omega*, 2023; 8(21): 19194–19211.
 19. Nematollahi MH, Mehrabani M, Hozhabri Y, Mirtajaddini M, Iravani S. Antiviral and antimicrobial applications of chalcones and their derivatives: from nature to greener synthesis. *Heliyon*, 2023; 9(10): e20428.
 20. Piekara J, Piasecka-Kwiatkowska D. Antioxidant potential of xanthohumol in disease prevention: evidence from human and animal studies. *Antioxidants*, 2024; 13(12): 1559.
 21. Rajendran G, Bhanu D, Aruchamy B, Ramani P, Pandurangan N, Bobba KN, Oh EJ, Chung HY, Gangadaran P, Ahn B. Chalcone: a promising bioactive scaffold in medicinal chemistry. *Pharmaceuticals*, 2022; 15(10): 1250.
 22. Kumari S, Sharma A, Yadav S. Novel coumarin chalcone derivatives: synthesis, docking and antimicrobial evaluation. *J Appl Pharm Res*, 2025; 13(1): 62–74.
 23. Yu M, Zhang X, Zhu H, Zhang X. Design, synthesis, molecular docking and anticancer activity of chalcone derivatives as VEGFR-2 inhibitors. *Molecules*, 2025; 30(23): 4526.
 24. Kachhi M. Synthesis and evaluative study of chalcones and their derivatives for antioxidant activity. *World J Pharm Res*, 2022; 11(8): 1–18.
 25. Patel RN, Patel PV. Synthesis and study of novel chalcone derivatives and their antimicrobial activity. *Eur J Exp Biol*, 2012; 2(5): 1492–1496.
 26. Dhanya O, Navya P, Mumthas RPV, Thasni MS, Saleem CKS. A review of microwave-assisted chalcone synthesis: advancements over conventional methods and their pharmacological actions. *Int J Pharm Sci Rev Res*, 2025; 85(3): 158–165.
 27. Rammohan A, Reddy JS, Sravya G, Rao CN, Zyryanov GV. Chalcone synthesis, properties and medicinal applications: a review. *Environ Chem Lett*, 2020; 18: 433–458.
 28. Lodhi GN, Nayak A, Jain AP. Chemical and pharmacological evolution of some synthesized chalcones and heterocyclic compounds. *Int J Adv Res Ideas Innov Technol*, 2021; 7(3): 2130–2140.
 29. Katade S, Phalgune U, Biswas S, Wakharkar R, Deshpande N. Microwave studies on synthesis of biologically active chalcone derivatives. *Indian J Chem B*, 2008; 47B: 927–931.
 30. Alam MS, Rahman SMM, Lee DU. Synthesis, biological evaluation, quantitative SAR and docking studies of novel chalcone derivatives as antibacterial and antioxidant agents. *Chem Pap*, 2015; 69(8): 1118–1129.
 31. Ali AM, Salah H, Gad MA, Youssef MAM, Elkanzi NAA. Design, synthesis and SAR studies of novel chalcone derivatives for potential insecticidal bioefficacy screening on *Spodoptera frugiperda*. *ACS Omega*, 2022; 7: 40091–40097.
 32. Shntaif AH. Green synthesis of chalcones under microwave irradiation. *Int J ChemTech Res*, 2016; 9(2): 36–39.
 33. Paul R, Sen S, Adak D, Porey A, Das S. A review: green synthesis of chalcones. *J Propuls Technol*, 2023; 44(6): 1534–1543.
 34. Bhuiyan MMH, Hossain MI, Mahmud MM, Al-Amin M. Microwave-assisted efficient synthesis of chalcones as probes for antimicrobial activities. *Chem J*, 2011; 1(1): 21–28.
 35. Kakati D, Sarma JC. Microwave-assisted solvent-free synthesis of 1,3-diphenylpropenones. *Chem Cent J*, 2011; 5(1): 8.
 36. Srivastava YK. Ecofriendly microwave-assisted synthesis of some chalcones. *Rasayan J Chem*, 2008; 1(4): 884–886.
 37. Gomes MN, Muratov EN, Pereira M, Peixoto JC, Rosseto LP, Cravo PVL, Andrade CH, Neves BJ. Chalcone derivatives: promising starting points for drug design. *Molecules*, 2017; 22(8): 1210.
 38. Dhaliwal JS, Moshawih S, Goh KW, Loy MJ, Hossain MS, Hermansyah A, Kotra V, Kifli N, Goh HP, Dhaliwal SKS, et al. Pharmacotherapeutic applications and chemistry of chalcone derivatives. *Molecules*, 2022; 27(20): 7062.
 39. Jayakumar A, Manikandan S, Mahalakshmi T, Kumar SM, Kumar BM. Microwave-assisted synthesized novel chalcone derivatives and evaluation of their antioxidant activity. *Indo Am J Pharm Sci*, 2023; 10(12): 519–524.
 40. Kumari A, Kumar S. Molecular docking studies and in silico ADMET screening of new chalcones as lanosterol α -demethylase and DNA gyrase inhibitors. *Orient J Chem*, 2025; 41(4): 1255–1268.
 41. Okolo EN, Ugwu DI, Ezema BE, Ndefo JC, Eze FU, Ezema CG, Ezugwu JA, Ujam OT. New chalcone derivatives as potential antimicrobial and antioxidant agents. *Sci Rep*, 2021; 11: 21781.
 42. Parmar I, Patel S, Shah U, Patel C, Patel A, Patel A. Synthesis, characterization, biological screening and in silico studies of novel pyrazole–chalcone derivatives as potent aromatase inhibitors for breast cancer therapy. *Future J Pharm Sci*, 2025; 11: 161.
 43. Ahmad MR, Sastry VG, Bano N, Anwar S. Synthesis of novel chalcone derivatives by conventional and microwave irradiation methods and their pharmacological activities. *Arab J Chem*, 2016; 9: S931–S935.
 44. Elkanzi NAA, Hrichi H, Alolayan RA, Derafa W, Zahou FM, Bakr RB. Synthesis of chalcone derivatives and their biological activities: a review. *ACS Omega*, 2022; 7(32): 27769–27786.
 45. Bukhari SNA, Jasamai M, Jantan I, Ahmad W. Review of methods and various catalysts used for

- chalcone synthesis. *Mini Rev Org Chem*, 2013; 10(1): 73–83.
46. Vogel AI. *Textbook of Practical Organic Chemistry*. 5th ed. London: Longman; 1989.
47. Nowakowska Z. A review of anti-infective and anti-inflammatory chalcones. *Eur J Med Chem*, 2007; 42(2): 125–137.
48. Kumar D, Rane N. Chalcone derivatives: synthesis and biological activities. *Mini Rev Med Chem*, 2013; 13(5): 596–604.
49. Sharma V, Sharma S, Prasad DN. Synthesis and characterization of chalcone derivatives. *Int J Chem Sci*, 2017.
50. Singh R, Prakash P. Synthesis and antimicrobial activity of chalcone derivatives. *Indian J Pharm Sci*, 2010; 72(2): 235–239.
51. Coates J. Interpretation of infrared spectra: a practical approach. In: *Encyclopedia of Analytical Chemistry*, 2000.
52. Pavia DL, Lampman GM, Kriz GS. *Introduction to Spectroscopy*. 5th ed. Cengage Learning; 2014.
53. LibreTexts Chemistry. Infrared spectroscopy.
54. Socrates G. *Infrared and Raman Characteristic Group Frequencies*. Wiley; 2001.
55. Balouiri M, Sadiki M, Ibsouda SK. Methods for in vitro evaluating antimicrobial activity: a review. *J Pharm Anal*, 2016; 6(2): 71–79.
56. Perez C, Pauli M, Bazerque P. An antibiotic assay by the agar well diffusion method. *Acta Biol Med Exp*. 1990.
57. Bonev B, Hooper J, Parisot J. Principles of assessing bacterial susceptibility. *J Antimicrob Chemother*; 2008; 61(6): 1295–1301.
58. CLSI. *Performance standards for antimicrobial susceptibility testing*. CLSI guideline M100, 2020.
59. Sakat SS, Juvekar AR. In vitro antimicrobial activity of medicinal compounds. *J Pharm Pharmacol*, 2010.
60. Trott O, Olson AJ. AutoDock Vina. *J Comput Chem*, 2010; 31(2): 455–461.
61. Morris GM, Huey R, Lindstrom W. AutoDock4. *J Comput Chem*, 2009; 30(16): 2785–2791.
62. Daina A, Michielin O, Zoete V. SwissADME. *Sci Rep*, 2017; 7: 42717.
63. BIOVIA DS. Discovery Studio Visualizer. Dassault Systèmes; 2019.
64. Berman HM, Westbrook J, Feng Z, et al. The Protein Data Bank. *Nucleic Acids Res*, 2000; 28(1): 235–242.
65. Lawson CL, et al. Protein Data Bank update. *Nucleic Acids Res*, 2011; 39: D456–D464.
66. Batovska DI, Todorova IT. Trends in utilization of chalcones as antimicrobial agents. *Curr Clin Pharmacol*, 2010; 5(1): 1–29.
67. Yadav VR, Prasad S. Chalcones: synthesis and biological activities. *Pharm Res*, 2011.
68. Go ML, Wu X, Liu XL. Chalcones: cytotoxic and antimicrobial properties. *Curr Med Chem*, 2005; 12(4): 483–499.
69. Bhat BA, Dhar KL. *Bioorg Med Chem Lett*, 2005; 15: 3177–3180.
70. Kontogiorgis CA, Hadjipavlou-Litina DJ. *J Med Chem*, 2005; 48: 6400–6408.