

**MANAGEMENT OF KARNINI YONIVYAPAD WITH APAMARGA KSHARA: A
CONCEPTUAL AYURVEDIC REVIEW****Dr. Sanju Lata Patel^{*1}, Dr. Shashi Sharma², Dr. Shikha Sharma³**¹P.G Scholar, ²Prof. & HOD, ³Lecturer,^{1,2,3}Department of Prasuti Tantra & Stri Rog, State Ayurvedic College & Hospital Lucknow, Maha Yogi Guru Gorakhnath Ayush University, Uttar Pradesh, India.***Corresponding Author: Dr. Sanju Lata Patel**

P.G Scholar, Department of Prasuti Tantra & Stri Rog, State Ayurvedic College & Hospital Lucknow, Maha Yogi Guru Gorakhnath Ayush University, Uttar Pradesh, India.

DOI: <https://doi.org/10.5281/zenodo.22159433>**How to cite this Article:** Dr. Sanju Lata Patel^{*1}, Dr. Shashi Sharma², Dr. Shikha Sharma³ (2026). Management Of Karnini Yonivyapad With Apamarga Kshara: A Conceptual Ayurvedic Review. World Journal of Pharmaceutical and Medical Research, 12(9), 44-47.

This work is licensed under Creative Commons Attribution 4.0 International license.

Article Received on 18/07/2026

Article Revised on 08/08/2026

Article Published on 01/09/2026

ABSTRACT

Karnini Yonivyapad is one of the twenty Yonivyapad described in Ayurvedic classics, characterized by the development of a Karnika-like growth at the Garbhashaya-dwara-mukha (cervix). Clinically, it presents with symptoms such as excessive mucoid vaginal discharge, itching, pelvic discomfort, and occasional bleeding. On the basis of morphological and symptomatic similarity, it can be correlated with cervical erosion (ectopy), a common gynecological condition in women of reproductive age. The present study is a conceptual review aimed at understanding the etiopathogenesis of Karnini Yonivyapad and evaluating the therapeutic potential of Apamarga Kshara in its management. According to Ayurveda, the disease is predominantly a Vata-Kapha disorder with Rakta involvement, leading to the formation of a localized granthi-like lesion. Apamarga Kshara, possessing Lekhana, Chedana, and Shodhana properties, is considered effective in removing abnormal tissue, reducing Kapha dominance, and promoting healing. This review highlights that Apamarga Kshara may act as an Ayurvedic counterpart to chemical cauterization used in modern medicine. It offers a minimally invasive, cost-effective approach for symptomatic cases. However, proper diagnosis and exclusion of malignancy are essential before intervention. Further clinical studies are needed to validate its efficacy and safety.

KEYWORDS: Karnini Yonivyapad, cervical erosion, cervical ectopy, Apamarga Kshara, Kshara Karma, Ayurveda.**INTRODUCTION**

Health of a woman has wide implications for family and society, and disorders of the reproductive tract therefore deserve careful attention. In Ayurveda, all diseases of the female genital tract are grouped under Yonivyapad, and Karnini Yonivyapad is one of the important entities described in this category. Classical reviews describe it as a disease in which a Karnika-like structure develops at the cervical opening, producing obstruction, discharge, and other symptoms that resemble cervical erosion in modern parlance.

The concept has been interpreted by recent authors as a condition that may correlate with cervical ectopy or erosion, especially because the cervix may appear red, granular, mildly hypertrophied, and covered with mucoid discharge. Associated symptoms such as pruritus,

backache, pelvic discomfort, contact bleeding, and infertility have also been discussed in these reviews.

From the Ayurvedic perspective, the pathogenesis involves vitiation of Vata and Kapha with Rakta participation, leading to the formation of a localized growth or granthi-like structure at the cervix. The therapeutic challenge is to remove or reduce the abnormal local lesion while restoring normal tissue integrity. In this context, Apamarga Kshara becomes clinically relevant because Kshara Karma is traditionally regarded as a powerful local measure for abnormal growths and chronic localized lesions.

AIM

To critically evaluate the concept of Karnini Yonivyapad in Ayurveda and to explore the therapeutic role of

Apamarga Kshara in its management with reference to cervical erosion.

OBJECTIVES

1. To review the classical Ayurvedic literature regarding Karnini Yonivyapad, including its Nidana, Samprapti, Rupa, and Chikitsa.
2. To establish a conceptual correlation between Karnini Yonivyapad and cervical erosion based on clinical features and pathology.
3. To analyze the pharmacological and therapeutic properties of Apamarga Kshara in the context of Kshara Karma.
4. To evaluate the mode of action of Apamarga Kshara in managing Karnini Yonivyapad from an Ayurvedic perspective.
5. To compare the Ayurvedic management approach with modern treatment modalities for cervical erosion.
6. To assess the advantages and limitations of Apamarga Kshara as a treatment modality.
7. To highlight the scope for future clinical research in the management of Karnini Yonivyapad using Apamarga Kshara.

MATERIALS AND METHODS

This is a conceptual narrative review. Ayurvedic classical sources describing Yonivyapad, Karnini, and Kshara Karma were reviewed, along with standard gynecological texts on cervical erosion/ectopy. The available journal reviews on Karnini Yonivyapad were also examined to understand the clinical correlation and treatment logic. The review focused on four domains: nidana, samprapti, clinical features, and therapeutic rationale of Apamarga Kshara.

No primary clinical data were collected. The article is intended as a theoretical synthesis for academic discussion and thesis writing.

RESULTS

The conceptual review of classical Ayurvedic literature and modern gynecological understanding reveals a strong correlation between Karnini Yonivyapad and cervical erosion/ectopy. The analysis highlights the following major findings.

1. Dosha-Dushya Involvement

Karnini Yonivyapad is predominantly a Vata-Kapha dominant disorder with involvement of Rakta Dhatu. Vata contributes to abnormal movement and structural distortion at the Garbhashaya-dwara, while Kapha leads to excessive mucous secretion, local heaviness, and growth formation. Rakta involvement explains the reddish appearance and occasional bleeding tendencies.

2. Structural and Morphological Correlation

The classical description of “Karnika” as a rounded, elevated, granthi-like structure resembles.

- Papillary or follicular cervical erosion.

- Nabothian follicles.

- Granular transformation zone of the cervix.

Modern descriptions of cervical erosion also mention a bright red, velvety, granular area surrounding the external os, which closely aligns with the Ayurvedic description.

3. Symptomatic Similarity

Both conditions share common clinical features.

- Profuse mucoid vaginal discharge
- Pruritus vulvae
- Lower abdominal pain and backache
- Occasional contact bleeding
- Infertility in chronic cases

These overlapping features strengthen the conceptual correlation.

4. Therapeutic Rationale of Apamarga Kshara

Apamarga Kshara demonstrates the following theoretical actions.

- **Lekhana (scraping):** Helps in removal of abnormal epithelial overgrowth
- **Chedana (excision-like effect):** Reduces localized granthi or Karnika
- **Shodhana (cleansing):** Eliminates Kapha-dominant secretions
- **Ropana (healing):** Promotes regeneration of normal epithelium

This aligns with the modern objective of cervical erosion treatment, which aims to destroy abnormal columnar epithelium and allow squamous regeneration.

5. Conceptual Clinical Outcome (Theoretical)

Based on Ayurvedic principles and analogous modern interventions.

- Reduction in discharge
- Decrease in cervical congestion
- Flattening of granular lesions
- Improved cervical integrity
- Possible improvement in fertility (in selected cases)

DISCUSSION

Karnini Yonivyapad is described in Ayurvedic classics as a disorder characterized by the formation of a Karnika-like structure at the Garbhashaya-dwara-mukha. The present conceptual analysis supports its close correlation with cervical erosion (ectopy), primarily based on similarities in site, morphology, and symptomatology. The classical description of a raised, granular, and obstructive lesion corresponds well with the modern depiction of a bright red, velvety area around the cervical os caused by columnar epithelial extension over the ectocervix.

From an Ayurvedic perspective, the pathogenesis of Karnini Yonivyapad involves vitiation of Vata and Kapha along with Rakta dushti. Vata contributes to structural derangement and displacement of tissue, while Kapha is responsible for excessive secretion, heaviness, and

growth formation. Rakta involvement explains the hyperemic appearance and occasional bleeding. These pathological changes collectively lead to the formation of Karnika, which can be interpreted as a granthi-like lesion at the cervix.

The clinical manifestations described in Ayurveda, such as Pichhila Yoni (mucoid discharge), Kandu (itching), Shoola (pain), and backache, show strong resemblance to the symptoms of cervical erosion, including vaginal discharge, pruritus, pelvic discomfort, and contact bleeding. This symptomatic overlap strengthens the conceptual bridge between the two conditions and justifies the application of Ayurvedic treatment principles in such cases.

Apamarga Kshara emerges as a rational therapeutic option in this condition due to its classical properties of Lekhana (scraping), Chedana (excision-like action), and Shodhana (cleansing). In Karnini Yonivyapad, where Kapha dominance and localized abnormal tissue growth are evident, Kshara acts by removing unhealthy tissue, reducing excessive secretion, and facilitating proper healing. Its action can be compared to the modern concept of chemical cauterization, where abnormal epithelial tissue is destroyed to allow regeneration of normal squamous epithelium.

Furthermore, Apamarga Kshara not only acts locally but also aligns with the Ayurvedic principle of Dosha

management. By reducing Kapha and clearing Srotorodha (obstruction), it helps restore the physiological environment of the cervix. This dual action—local and systemic—gives it an advantage over purely mechanical or physical modern interventions.

In comparison with modern treatments such as electrocautery, cryotherapy, and laser ablation, Apamarga Kshara can be considered a minimally invasive, cost-effective, and Ayurveda-based alternative. While modern techniques focus mainly on tissue destruction, Kshara therapy integrates tissue removal with doshic balance, which may contribute to better symptomatic relief and reduced recurrence.

However, it is important to emphasize that proper case selection is crucial. Cervical erosion may mimic premalignant or malignant conditions; therefore, diagnostic procedures such as Pap smear and biopsy should be performed before initiating Kshara therapy. This ensures safety and prevents mismanagement of serious conditions.

Overall, Apamarga Kshara represents a promising Ayurvedic approach for the management of Karnini Yonivyapad, especially in symptomatic and benign cases. Nevertheless, further clinical studies and standardization protocols are required to establish its efficacy and safety on a broader scientific basis.

Advantages and Limitations of Apamarga Kshara in Karnini Yonivyapad

Table No- 1.

Aspect	Advantages of Apamarga Kshara	Limitations of Apamarga Kshara
Therapeutic Action	Provides Lekhana, Chedana, and Shodhana action on abnormal tissue	Depth of action is difficult to standardize
Mode of Action	Acts locally and helps in removal of pathological tissue	Risk of over-application leading to tissue damage
Dosha Effect	Effective in Kapha-dominant conditions and clears Srotorodha	Limited effect if Pitta predominance or acute inflammation is present
Healing	Promotes regeneration of healthy epithelium (Ropana)	Healing time may vary depending on patient condition
Cost & Accessibility	Cost-effective and easily available Ayurvedic intervention	Requires trained practitioner for proper application
Comparison with Modern Methods	Comparable to chemical cauterization; minimally invasive	Lacks large-scale clinical trial validation
Recurrence	May reduce recurrence by correcting Dosha imbalance	Recurrence possible if Nidana is not avoided
Safety	Generally safe when properly indicated and applied	Unsafe without proper diagnosis (e.g., malignancy not ruled out)
Integration	Can be combined with internal Ayurvedic medicines	Limited acceptance in modern clinical practice
Patient Compliance	Simple procedure, no major surgical setup required	May cause local discomfort or burning sensation

CONCLUSION

Karnini Yonivyapad can be conceptually correlated with cervical erosion or ectopy on the basis of clinical and morphological similarity. Apamarga Kshara is a rational Ayurvedic local therapeutic option in symptomatic,

benign, and properly investigated cases because of its Lekhana, Chedana, and Shodhana properties. Its use should be guided by careful gynecological evaluation and should always exclude premalignant or malignant cervical disease before treatment.

REFERENCES

1. Agnivesha. *Charaka Samhita*. Part 2. Revised by Charaka and Dridhabala, edited by Satyanarayana Shastri, Kasinath Shastri, Gorakhnath Chaturvedi, Rajeswaradatta Shastri. Varanasi: Chaukhambha Bharati Academy; Reprint 2006. Chikitsa Sthana, Chapter 30, Yonivyapachikitsitam.
2. Sushruta. *Sushruta Samhita*. Part 2. Edited by Ambikadutta Shastri. Varanasi: Chaukhambha Sanskrit Sansthan; Reprint 2006. Uttaratantra, Chapter 38, Yonivyapatpratishedha.
3. Vagbhata. *Ashtanga Sangraha*. Edited by S.P. Sharma. Varanasi: Chaukhambha Sanskrit Series Office; 2006. Uttara Tantra, Chapter 38.
4. Vagbhata. *Ashtanga Hridayam*. Edited by Atrideva Gupta. Varanasi: Chaukhambha Prakashan; Reprint 2008. Uttara Tantra, Chapter 33.
5. Vriddha Jivaka. *Kashyapa Samhita*. Varanasi: Chaukhamba Visvabharati Bhavan; Reprint 2013. Siddhi Sthana, Chapter 4.
6. Bhavamishra. *Bhavaprakasha Samhita*. Varanasi: Chaukhambha Sanskrit Bhavan; edition details as per standard text.
7. Narahari. *Yogaratanakara*. Varanasi: Chaukhambha Sanskrit Sansthan; standard edition.
8. Sushruta. *Sushruta Samhita*. Edited with Ayurveda-Tattva-Sandipika by Ambikadutta Shastri. Varanasi: Chaukhambha Sanskrit Sansthan; Reprint 2007. Sutra Sthana, Chapter 11, Kshara Agnikarma Vidhi.
9. Pal S, Prakash C. A critical review on Karnini Yonivyapad w.s.r. to cervical erosion. *Ayushdhara*, 2017; 4(6): 1478-1485.
10. Sushma D. Diagnosis and management of Karnini Yonivyapad w.s.r. to Cervical Erosion and its significance in Modern Era. *Journal of Ayurveda and Integrated Medical Sciences*, 2021; 6(4): 303-310.
11. Dutta DC. *Textbook of Gynecology*. New Delhi: Jaypee Brothers Medical Publishers; standard edition.
12. Berek JS, editor. *Berek & Novak's Gynecology*. Philadelphia: Wolters Kluwer; standard edition.
13. Shaw RW, Soutter WP, Stanton SL. *Gynaecology*. Edinburgh: Churchill Livingstone; standard edition.
14. Konar H. *DC Dutta's Textbook of Gynecology*. New Delhi: Jaypee Brothers Medical Publishers; standard edition.
15. The Ayurvedic Pharmacopoeia of India. Part I, Volume 1. New Delhi: Government of India, Ministry of AYUSH; standard edition.
16. Kirtikar KR, Basu BD. *Indian Medicinal Plants*. Dehradun: International Book Distributors; standard edition.
17. Nadkarni KM. *Indian Materia Medica*. Mumbai: Popular Prakashan; standard edition.
18. Sharma PV. *Dravyaguna Vijnana*. Varanasi: Chaukhambha Bharati Academy; standard edition.
19. Guleria K, et al. Classical concepts of Kshara Karma in anorectal and gynecological disorders. *Journal of Ayurveda and Integrated Medical Sciences*. standard citation format as applicable.
20. Sharma A, et al. Local Ayurvedic management of cervical ectopy: a conceptual review. *International Journal of AYUSH and Allied Systems*. standard citation format as applicable.