

**ROLE OF CHASE AMPUTATION AFTER INDEX FINGER TRAUMA : ABOUT ONE
CASE AND REVIEW OF LITERATURE**

***M. EL Mhadder, M. Abakka, R. Boueld, H. Berrada, Y. Mokhchani, A. Bennis, O. Zadoug, M. Benchakroun
and S. Bouabid**

Orthopedic Trauma Department 1, Mohammed V Military Training Hospital, Rabat.

***Corresponding Author: M. EL Mhadder**

Orthopedic Trauma Department 1, Mohammed V Military Training Hospital, Rabat.

Article Received on 01/07/2023

Article Revised on 21/07/2023

Article Accepted on 11/08/2023

ABSTRACT

Chase's intervention expands the treatment objectives by indexing the middle finger, which will ensure the pollicidigital grip, retain strength and respect the aesthetic appearance of the hand. It is a frequently performed surgical intervention, which gives very good results for the patients concerned. This study relates our experience in the Traumatology and Orthopedics Department of the Military Hospital in Rabat in the treatment of severe trauma of the index finger.

KEYWORDS: Chase-Index-Finger-Amputation.

1. INTRODUCTION

Chase's intervention extends to the treatment objectives by indexing the middle finger which will ensure the pollicidigital grip; maintain strength and respect the aesthetic appearance of the hand.

The Chase amputation is an amputation at the base of the second metacarpal, which allows indexing of the middle finger. It has the advantage of ensuring maximum opening of the middle thumb commissure.

Indexing the middle finger leads to better integration and usefulness of the finger.

We report the case of a 40-year-old patient,^[8] a patient with no previous history suffered a traumatic amputation of the index finger after handling a mold. The aesthetic and functional result was very satisfactory after six months of the operation.^[7]

2. RESEARCH METHODS

This is Mr. yassine E, a 40-year-old patient,^[7] with no previous history, who presented to the emergency room of the Military Hospital in Rabat for a traumatic amputation of the index finger after handling a mold,^[7] The patient was initially trimmed surgically with a compressive dressing, then admitted to the operating room for regularization using the Chase technique.

3. RESULTS

The patient underwent amputation using the Chase technique, which consists of an amputation at the base of

the second metacarpal.^[4] allowing indexing of the middle finger. He also benefited from early rehabilitation.^[5] The control after 6 months shows a good aesthetic and functional result of the hand with a good opening of the thumb-medius commissure, indexing of the middle finger with patient satisfaction concerning the aesthetic and functional aspect of his hand.



Pictures before and after chase's amputation.

4. DISCUSSION

The index finger is the most used finger after the thumb.^[7] It is an essential element of the pollicidigital grip and a stabilizing element in the overall digitopalmar grip.

The Chase amputation is an amputation at the base of the second metacarpal, which allows the indexing of the middle finger,^[5] it has the advantage of ensuring maximum opening of the thumb-medius commissure. The operating protocol includes two essential steps.^[6]

- Cutaneous time: the incision is most often dorsal in order to avoid a potentially embarrassing palmar scar, it circumscribes the base of the index finger like a “snowshoe”;
- Tendon and bone time: by dorsal approach, the extensor apparatus is sectioned. The extensor indicis tendon is cut above the metacarpophalangeal joint, then transferred to the extensor digitorum tendon intended for the middle finger. The flexor tendons are sectioned proximally, wrist in flexion.
- Vasculonervous time: the collateral nerves, dissected proximally, are sectioned high in the palm of the hand then hemostasis is ensured.

5. CONCLUSION

Surgery is the standard treatment for traumatic amputation of the index finger.^[4] However, the function of the hand must also be taken into consideration; the principle is not to maintain a level that is not functionally better than that which would have been obtained by shortening the bone; to give correct hand function.^[3]

REFERENCES

1. Kapandji, A. I. (January). La préhension dans la main humaine. In *Annales de Chirurgie de la Main* (Vol. 8, No. 3, pp. 234-241). Elsevier Masson, 1989.
2. Littler, J. W. Architectural principles of reconstructive hand: Surgery. *Surg Clin North Am*, 1951; 31(2): 463-476.
3. Chase, R. A. Surgery of the hand. *N Engl J Med*, 1972; 287(23): 1174-1181.
4. Chase, R. A. The damaged index digit: a source of components to restore the crippled hand. *JBJS*, 1968; 50(6): 1152-1160.
5. Erhard, L., Medina, J., Zabo, S., Pajardi, G., & Foucher, G. (February). Secondary treatment of digital mutilations: reconstruction or amputation. In *Annales de chirurgie plastique et esthetique*, 2002; 47(1): 47-56.

6. Murray, J. F., Carman, W., & MacKenzie, J. K. Transmetacarpal amputation of the index finger: a clinical assessment of hand strength and complications. *The Journal of hand surgery*, 1977; 2(6): 471-481.
7. Archives of the Traumatology-Orthopedics Department of the CHU Ibn Sina in Rabat.
8. Iconography of the Traumatology-Orthopedics duty team of the Ibn Sina Hospital in Rabat.
9. EM Consulte: Article on amputation and disarticulation of limbs.