

EFFECT OF VITAMIN D SUPPLEMENTATION IN BRONCHIAL ASTHMA

*¹Dr. Saira Gulzar, ²Dr. Hajra Arif and ³Dr. Saher Iftikhar¹PMDC # 89542-P.²PMDC # 88037-P.³PMDC # 87792-P.

*Corresponding Author: Dr. Saira Gulzar

PMDC # 89542-P

Article Received on 19/05/2018

Article Revised on 09/06/2018

Article Accepted on 30/06/2018

ABSTRACT

Background and Objective: Vitamin D deficiency has strong association with various respiratory disorders in which bronchial asthma is one of them. The objective was to determine the efficacy of vitamin D supplementation in cases with bronchial asthma. **Methods:** This case control study was conducted at Allama Iqbal memorial hospital, Sialkot from August to October 2017 in which 100 cases of bronchial asthma were randomly divided into Group-A and Group-B each contained 50 patients. Group-A was given placebo and Group-B with vitamin D in a dose of 50,000 units per day orally. Both the groups were followed in terms of improvement in FEV1 at 1, 2 and 3 months. **Results:** There was no significant difference in both groups in terms of BMI and duration of asthma at start of study. The mean pre treatment vitamin D level of Group-A was 14.23 ± 1.66 and of Group-B, 15.30 ± 2.05 ng/dl ($p = 0.23$). FEV1 in pre treatment Group-A was 64.35 ± 3.16 and of Group-B was 62.35 ± 2.16 with $p = 0.95$. There was no significant difference in terms of FEV1 in both the groups at one month ($p = 0.32$). While at two months it was significantly higher in Group-B with $p = 0.04$. At 3 months the final outcome was seen where the post treatment FEV1 in Group-A was 66.13 ± 2.75 and in Group-B, 75.15 ± 2.04 with p value of 0.001. **Conclusion:** Vitamin D supplementation improves FEV1 significantly at two months and these can be even highly significant if it is extended up to 3 months.

KEYWORDS: Vitamin D, asthma.

INTRODUCTION

Asthma is one of the major health issues across the world from childhood to adult life. According to a survey more than 25 million people are suffering from this in United States in 2012 and is one of the most reported entity in both outpatient and inpatient departments. It posed a great burden in terms of economy. The role of environmental factors in the prevalence of asthma cannot

be denied but recent researches have focused more on metabolic and nutritional factors in which vitamin D level is one of them. Despite all the advances in the medical sciences and researches there are no definitive treatment to eradicate it and the options used are to control it and sometimes fail to do it, for which newer agents are always in need.^[1,2]

Table I: Comparison between variables of two groups.

Variable	Group-A	Group-B	p-value
Male/Female	26/24	29/21	0.87
Age (years)	26.50 ± 5.50	27.60 ± 6.50	1.04
BMI (Kg)	24.75 ± 3.20	26.70 ± 6.40	0.96
Duration of asthma	12.40 ± 3.50	11.66 ± 3.05	0.66
Pre-treatment vit D level	14.23 ± 1.66	15.30 ± 2.05	0.23
Pre treatment FEV 1	64.35 ± 3.16	62.35 ± 2.16	0.95

Vitamin D is one of the newer agents that are being studied in the treatment of such cases. Vitamin D has been thought to act in various ways. Alpha hydroxylase is present in the lung epithelium which is capable of converting calcidiol to calcitriol; and this is thought to

interfere in the synthesis of different cytokine i.e. RANTES, PDGF, interleukin, different CD cells and metalloproteinases. These are the markers of inflammation found in the cases with asthma and enhance the inflammatory mechanisms. Vitamin D is

presumed to decrease in the infection rate as well due to cathelicidin, which is over expressed with its supplementation and is anti infective in nature.^[3,4]

Extensive work up is being done in recent times to build the association of its level with various respiratory conditions and in many circumstances it is also proved that the low levels of 25-hydroxyvitamin D are associated with increased respiratory tract infection, decreased lung function tests, reduced steroid response and frequent exacerbation.^[5-8]

The present study was designed to see the effect of vitamin D supplementation in the improvement of symptoms of bronchial asthma in terms of FEV1.

METHODS

This case control study was conducted at Allama Iqbal memorial Hospital, Sialkot from August to October 2017 in which 100 cases of bronchial asthma aged 18 to 50 years were randomly divided into Group-A and Group-B each contained 50 patients. The diagnosis of bronchial asthma was made by PFTs with FEV1/FVC ratio of 0.7 and reversibility in FEV 1 of more than 12% in cases with history of cough, shortness of breath or chest tightness. The cases with history of renal, liver disease, pregnancy were excluded from the study. The demographics like age, BMI, duration of asthma and

baseline level of vitamin D was checked and data was collected. The cases were asked to select one of the shield opaque envelope labeled as A or B. The cases in Group-A were treated as per GINA guidelines 2016 according to severity of disease along with placebo and Group-B was also treated with anti-asthmatic drugs as per severity and vitamin D in a dose of 50,000 units per day orally. Both the Groups were followed in terms of FEV1 at one month, two months and three months where final outcome was seen. Efficacy was assessed in terms of improvement in FEV1.

Statistical Analysis: A sample of 50 cases in each group was selected. Both the groups were compared in terms of their mean age, BMI, vitamin D level, baseline FEV1. Data was stratified to see for significance in both groups regarding efficacy. Independent sample t-test and chi square test were used to compare the both groups taking p-value of ≤ 0.05 as significant.

RESULTS

There were total 100 cases, 50 in each group. Group-A contained 26 (52%) males and 24 (48%) females, and Group-B has 29 (58%) males and 21 (42%) females (Table-I). The mean age of Group-A was 26.50 ± 5.50 and Group-B, 27.60 ± 6.50 years with $p=1.04$. There was no significant difference in both groups in terms of BMI and duration of asthma.

Table II: Efficacy comparison between two groups.

FEVI 1	Group-A	Group-B	Significance
FEV 1 at baseline	64.35 \pm 3.16	62.35 \pm 2.16	p= 0.95
FEV 1 at 1 month	65.65 \pm 3.05	66.28 \pm 1.10	p= 0.32
FEV 1 at 2 months	65.84 \pm 2.95	71.44 \pm 2.78	p= 0.04
FEV 1 at 3 months	66.13 \pm 2.75	75.15 \pm 2.04	p= 0.001

The mean pre treatment vitamin D level of Group-A was 14.23 ± 1.66 and of Group-B, 15.30 ± 2.05 ng/ dl ($p= 0.23$). FEV1 in pre treatment Group-A was 64.35 ± 3.16 and of Group-B was 62.35 ± 2.16 with $p= 0.95$ as in Table-I. There was no significant difference in terms of FEV1 in both the groups at one month ($p= 0.32$). While at two months it was significantly higher in Group-B with $p= 0.04$. At three months the final outcome was seen where the post treatment FEV1 in Group-A was 66.13 ± 2.75 and in Group-B, 75.15 ± 2.04 with p value of 0.001 as in Table-II.

DISCUSSION

Bronchial asthma is a high burden disease in developing countries like Pakistan and their management relies upon the pharmacotherapies for which new guidelines are recommended every year by NICE group and are tried to meet. But in some cases even with these controller and reliever medications, there are cases, which are still not responding to conventional treatments. This again calls for looking for the other factors, which can be associated with these, as asthma is multifactorial and different

pathophysiological mechanisms are involved with different degrees of severity in individual cases. Recent data also support that the current treatment options used in the treatment of asthma are not enough and newer and better options are required.

In this study there was significant difference in terms of FEV1 in both the groups at two months with $p= 0.04$. At three months the final outcome was seen where the post treatment FEV1 in Group-A 66.13 ± 2.75 and in Group-B, 75.15 ± 2.04 with p value of 0.001. This was also proved by other studies as well that the asthma symptoms were improved with treatment of vitamin D supplementation. In a study done by Black and Scragg et al., they found strong association in improvement of asthma symptoms in term of FEV 1 and improvement in FVC with vitamin D3.^[9] While in recent study done by Gargen et al. in 2013, they used vitamin D supplementation and reported that the cases treated with this have lower chance of respiratory tract infections especially allergic conditions and also revealed that the higher the concentration of vitamin D and lower are the levels seen of IgE level and peripheral eosinophilia.^[10] This again re enforces its use

in cases of asthma considering it as enhanced allergic phenomenon and those cases where there is increased eosinophil and IgE level.

Bergman *et al* also report decreased incidence of respiratory infections in vitamin D augmented group.^[11] However, studies done by Iqbal *et al.* also reported improvement in inflammation in cases with asthma, however they did not find any significant association with this.^[12]

In a study done by Dupont *et al.*^[13] they added leukotriene receptor blocker with vitamin D and study done by Korn *et al.*, Shah *et al* and Keith *et al* also used it as an add on and revealed improvement in cases with asthma.^[14-16] However they studies these population for 8 weeks at which we also found significant improvement with p-value of 0.04 but at 3 months we found even more significant improvement with p= 0.001.

CONCLUSION

Vitamin D supplementation improves FEV1 significantly at eight weeks and these can be even highly significant if it is extended up to three months.

REFERENCES

1. Asthma fast stats. Centers for disease control and prevention website. Accessed January 05, 2017 (internet). Available from; <http://www.cdc.gov/nchs/faststats/asthma.htm>.
2. Sears MR. Trends in the prevalence of asthma. *Chest Journal.*, 2014; 142(2): 219-225.
3. Ianagan JN, Wang L, Tangpricha V. Regulation of the 25-hydroxyvitamin D-1alpha-hydroxylase gene and its splice variant. *Recent Results Cancer Res.*, 2003; 164: 157-167.
4. Hewison M, Zehnder D, Bland R. 1 alpha-Hydroxylase and the action of vitamin D. *J Mol Endocrinol.*, 2000; 25: 141-148.
5. Sutherland ER, Goleva E, Jackson LP, Stevens AD, Leung DY. Vitamin D levels, lung function, and steroid response in adult asthma. *Am J Respir Crit Care Med.*, 2010; 181: 699-704.
6. Goleva E, Searing DA, Jackson LP, Richers BN, Leung DY. Steroid requirements and immune associations with vitamin D are stronger in children than adults with asthma. *J Allergy Clin Immunol.*, 2012; 129: 1243-1251.
7. Brehm JM, Celedon JC, Soto-Quiros ME. Serum vitamin D levels and markers of severity of childhood asthma in Costa Rica. *Am J Respir Crit Care Med.*, 2009; 179: 765-771.
8. Wu AC, Tantisira K, Li L, Fuhlbrigge AL, Weiss ST, Litonjua A. The effect of vitamin D and inhaled corticosteroid treatment on lung function in children. *Am J Respir Crit Care Med.*, 2012; 186: 508-513.
9. Black PN, Scragg R. Relationship between serum 25-hydroxyvitamin d and pulmonary function in the third national health and nutrition examination survey. *Chest.*, 2005; 128: 3792-3798.
10. Gergen PJ, Teach SJ, Mitchell HE, Freishtat RF, Calatroni A, Matsui E, *et al.* Lack of a relation between serum 25-hydroxyvitamin D concentrations and asthma in adolescents. *Am J Clin Nutr.*, 2013; 97: 1228-1234.
11. Bergman P, Norlin AC, Hansen S, Rekha RS, Agerberth B, Björkhem-Bergman L, *et al.* Vitamin D 3 supplementation in patients with frequent respiratory tract infections: a randomised and double-blind intervention study. *BMJ Open.*, 2012; 2: 1-10.
12. Muhammad Zafar Majeed Babar *et al* qbal SF, Freishtat RJ. Mechanism of action of Vitamin D in the asthmatic lung. *J Investig Med.*, 2011; 59: 12001202.
13. Dupont L, Potvin E, Korn D, Lachman A, Dramaix M, Gusman J, *et al.* Improving asthma control in patients suboptimally controlled on inhaled steroids and long-acting beta2-agonists: Addition of montelukast in an open-label pilot study. *Curr Med Res Opin.*, 2005; 21: 863-869.
14. Korn D, Van den Brande P, Potvin E, Dramaix M, Herbots E, Peché R. Efficacy of add-on montelukast in patients with non-controlled asthma: a Belgian open-label study. *Curr Med Res Opin.*, 2009; 25: 489-497.
15. Shah AR, Sharples LD, Solanki RN, Shah KV. Double-blind, randomised, controlled trial assessing controller medications in asthma. *Respiration.*, 2006; 73: 449-456.
16. Keith PK, Koch C, Djandji M, Bouchard J, Psaradellis E, Sampalis JS, *et al.* Montelukast as add-on therapy with inhaled corticosteroids alone or inhaled corticosteroids and long-acting beta-2-agonists in the management of patients diagnosed with asthma and concurrent allergic rhinitis (the RADAR trial). *Can Respir J.*, 2009; 16: 17-31.