

**ABDOMINAL LYMPH NODE METASTASIS OF BREAST CARCINOMA**

Dr. F. Z. Lahlali, H. Sfaoua\*, A. Allam, J. Aarab, S. El Majjaoui, H. El Kacemi, T. Kibdani, N. Benjaafar

National Institute of Oncology, Radiotherapy Department Mohammed 5 University, Rabat- Morocco.

\*Corresponding Author: H. Sfaoua

National Institute of Oncology, Radiotherapy Department Mohammed 5 University, Rabat- Morocco.


Article Received on 29/03/2019

Article Revised on 19/04/2019

Article Accepted on 09/05/2019

**Clinical image**

A 47-year-old female, G0P0, she had no history of oral contraception use, neither pathologic disease, who presented Invasive breast carcinoma of no special type, staged T2N1M0, with nodal extracapsular extension, SBR III, ER /PR positive. The patient undergone surgery, chemotherapy, radiation therapy and hormonal therapy. Follow up showed a good clinical response during eight years. Abdominal complaints caused the patient to consult, She proceeded to have a diagnostic abdominal scan, which demonstrated a huge abdominal lymph nodes, realizing intra and retroperitoneal masses. Biopsies revealed a metastasis of breast carcinoma.



CT scan showing of abdominal adenopathy realizing intra and retroperitoneal masses (A sagittal plane, B coronal plane, C axial plan).