

AN OVERVIEW ON ASHWAGANDHA: A RASAYANA (REJUVENATOR) OF  
AYURVEDA: A REVIEWDr. Vishesh Duggal<sup>\*1</sup>, Dr. Omprakash Sharma<sup>2</sup> and Dr. Naresh Garg<sup>2</sup><sup>1</sup>PG Scholar Department of Dravyaguna Vigyan.<sup>2</sup>Professor and HOD Department of Dravyaguna Vigyan, SriGanganagar College of Ayurvedic Science and Hospital, Tanta University, SriGanganagar – 335001, India.**\*Corresponding Author: Dr. Vishesh Duggal**  
PG Scholar Department of Dravyaguna Vigyan.

Article Received on 28/08/2021

Article Revised on 18/09/2021

Article Accepted on 08/10/2021

**ABSTRACT**

*Withania somnifera* (Ashwagandha) is very revered herb of the Indian Ayurvedic system of medicine as a Rasayana (tonic). It is used for various kinds of disease processes and specially as a nervine tonic. Considering these facts many scientific studies were carried out and its adaptogenic / anti-stress activities were studied in detail. In experimental models it increases the stamina of rats during swimming endurance test and prevented adrenal gland changes of ascorbic acid and cortisol content produce by swimming stress. Pretreatment with *Withania somnifera* (WS) showed significance protection against stress induced gastric ulcers. In some cases of uterine fibroids, dermatosarcoma, long term treatment with WS controlled the condition. It has a Cognition Promoting Effect and was useful in children with memory deficit and in old age people loss of memory. It was also found useful in neurodegenerative diseases such as Parkinson's, Huntington's and Alzheimer's diseases. It has GABA mimetic effect and was shown to promote formation of dendrites. It has anxiolytic effect and improves energy levels and mitochondrial health. It is an anti-inflammatory and anti-arthritis agent and was found useful in clinical cases of Rheumatoid and Osteoarthritis. Large scale studies are needed to prove its clinical efficacy in stress related disorders, neuronal disorders and cancers.

**KEYWORDS:** *Withania somnifera*, rejuvenator, adaptogen / anti-stress, anti-tumor, neuroregenerative, anti-arthritis.**INTRODUCTION**

Ashwagandha (*Withania somnifera*, fam. Solanaceae) is commonly known as “Indian Winter cherry” or “Indian Ginseng”. It is one of the most important herb of Ayurveda (the traditional system of medicine in India) used for millennia as a Rasayana for its wide ranging health benefits. Rasayana is described as an herbal or metallic preparation that promotes a youthful state of physical and mental health and expands happiness. These types of remedies are given to small children as tonics, and are also taken by the middle-aged and elderly to increase longevity. Among the ayurvedic Rasayana herbs, Ashwagandha holds the most prominent place. It is known as “Sattvic Kapha Rasayana” Herb. Most of the Rasayana herbs are adaptogen / anti-stress agents.

Ashwagandha is commonly available as a churna, a fine sieved powder that can be mixed with water, ghee (clarified butter) or honey. It enhances the function of the brain and nervous system and improves the memory. It improves the function of the reproductive system promoting a healthy sexual and reproductive balance. Being a powerful adaptogen, it enhances the body's

resilience to stress. Ashwagandha improves the body's defense against disease by improving the cell-mediated immunity. It also possesses potent antioxidant properties that help protect against cellular damage caused by free radicals.

**Chemical Composition**

The biologically active chemical constituents of *Withania somnifera* (WS) include alkaloids (isopelletierine, anaferine, cuseohygrine, anahygrine, etc.), steroidal lactones (withanolides, withaferins) and saponins. Saponins and acylsterylglucosides in Ashwagandha are anti-stress agents. Active principles of Ashwagandha, for instance the saponins VII-X and Withaferin-A,

**Classical Uses of Ashwagandha**

Ayurveda, the traditional system of medicine practiced in India can be traced back to 6000 BC (Charak Samhita, 1949). For most of these 6000 years Ashwagandha has been used as a Rasayana. The root of Ashwagandha is regarded as tonic, aphrodisiac, narcotic, diuretic, anthelmintic, astringent, thermogenic and stimulant. The

root smells like horse (“*ashwa*”), that is why it is called Ashwagandha (on consuming it gives the power of a horse). It is commonly used in emaciation of children (when given with milk, it is the best tonic for children), debility from old age, rheumatism, vitiated conditions of vata, leucoderma, constipation, insomnia, nervous breakdown, goiter etc. The paste formed when roots are crushed with water is applied to reduce the inflammation at the joints. It is also locally applied in carbuncles, ulcers and painful swellings. The root in combination with other drugs is prescribed for snake venom as well as

in scorpion-sting. It also helps in leucorrhoea, boils, pimples, flatulent colic, worms and piles. The Nagori Ashwagandha is the supreme among all Ashwagandha varieties. Maximum benefit appears when fresh Ashwagandha powder is used.

The leaves are bitter and are recommended in fever, painful swellings. The flowers are astringent, depurative, diuretic and aphrodisiac. The seeds are anthelmintic and combined with astringent and rock salt remove white spots from the cornea.



A

B

Images showing ashwagandha plant A & B.

#### Physiochemical & Pharmacological Effect

Ashwagandha is compared well with *Eleutherococcus senticosus* (Siberian Ginseng) and *Panax Ginseng* (Chinese / Korean Ginseng) in its adaptogenic properties, and hence it is popularly known as Indian Ginseng. The extensive studies on the biological model of animals for the adaptogenic / anti-stress properties of Ashwagandha have shown it to be effective in increasing the stamina (physical endurance) and preventing stress induced gastric ulcer, carbon tetrachloride (CCl<sub>4</sub>) induced hepatotoxicity and mortality. An aqueous suspension of Ashwagandha root was used at 100 mg/kg/oral dosage. The results indicate a significant increase in the plasma corticosterone level, phagocytic index and avidity index in rats subjected to cold swimming stress. In the rats pretreated with the drug, these parameters were near control values and an increase in the swimming time was observed. These results indicate that *Withania somnifera* used in the crude form is a potent anti-stress agent. The results of above studies lend support to the hypothesis of tonics, vitalizers and rejuvenators of Ayurveda which indicate clinical use of *Withania somnifera* in the prevention and treatment of many stress induced diseases like arteriosclerosis, premature ageing, arthritis, diabetes, hypertension and malignancy.

#### i. Effect on swimming performance

Ashwagandha was shown to increase swimming performance in rats as judged by increase in swimming time during physical endurance test. Ashwagandha's antistress properties have been investigated in all these studies using adult rats were carried out by swimming endurance stress test. Ashwagandha treated animals showed a significant increase in the duration of swimming time as compared to control. The control group of mice swam for a mean time of 385 minutes, whereas the drug-treated animals continued to swim for a mean duration of 740 minutes. Thus, the swimming time was approximately doubled after *Withania somnifera* (WS) treatment.

#### ii. Effect on cortisol and ascorbic acid contents of adrenals

The cortisol content of adrenals was reduced significantly in animals subjected to 5 h constant swimming as compared to non-swimmer group. Pretreatment with WS prevented reduction of the cortisol content of adrenals. The ascorbic acid content was also reduced significantly after 5 h of swimming as compared to the animal of non-swimmer group. Pretreatment with WS prevent reduction in ascorbic acid content which occurs after swimming stress. Thus, *Withania somnifera* treatment prevents, decrease of adrenal

cortisol and ascorbic acid which occurs due to swimming stress.

### iii. Anti-ulcerogenic effect

Ashwagandha was found to be useful in the prevention of stress-induced ulcers of the gastrointestinal tract. It showed significant protection against 18 h immobilization, cold + immobilization (4h) and aspirin induced gastric ulcers and lowered the mean ulcer index in rats.

### iv. Effect on leucocytosis

Ashwagandha given to a group of mice with milk injection produced reduction in leucocytosis.

### v. Anabolic effects

#### Effect on Central Nervous System Cognition Promoting Effect

Ashwagandha is a well-known Ayurvedic Rasayana, and belongs to a sub-group of Rasayanas known as Medhyarasayanas. Medhya typically refers to the mind and mental/intellectual capacity. Thus, Medhya Rasayana like Ashwagandha, is used to promote intellect and memory. The cognition-promoting effect of Medhya Rasayanas is best seen in children with memory deficits, or when memory is compromised following head injury, or a prolonged illness and in old age.

#### Effect on neurodegenerative diseases such as Parkinson's, Huntington's and Alzheimer's diseases

In patients with Alzheimer's disease, neuritic atrophy and synaptic loss are considered the major causes of cognitive impairment, as based on the results of neuropathological post-mortem studies of the brain. In the brains of patients suffering from other neurodegenerative diseases such as Parkinson's disease, Huntington's disease, and Creutzfeldt- Jakob disease, the atrophy of neurites has also been observed as a significant part of the etiology. There are dozens of studies that show that Ashwagandha slows, stops, reverses or removes neuritic atrophy and synaptic loss. Therefore, Ashwagandha can be used to treat Alzheimer's, Parkinson's, Huntington's and other neurodegenerative diseases at any stage of the disease, even before a person has been diagnosed and is still in the state of mild forgetfulness, etc. Glycowithanolides withaferin- A and sitoindosides VII-X isolated from the roots of Ashwagandha significantly reversed ibotenic acid induced cognitive defects in Alzheimer's disease model.

Ashwagandha has been described as a nervine tonic in Ayurveda and that is why it is a common ingredient of Ayurvedic tonic. Tonics, rejuvenators and vitalizers of Ayurveda appear to allay disease and induce immunity and longevity in the users.

Pretreatment with Ashwagandha extract was found to prevent all the changes in antioxidant enzyme activities, catecholamine content, dopaminergic D2 receptor

binding and tyrosine hydroxylase expression induced by 6-hydroxydopamine (6-OHDA) in rats (an animal model of Parkinson's disease) in a dose-dependent manner. Thus, these results suggest that Ashwagandha may be helpful in protecting the neuronal injury in Parkinson's disease.

#### GABA-mimetic effect on neurodegeneration and neuroregenerative potential

Behavioral experiments have lent support to the GABA-mimetic activity of Ashwagandha root extract. GABAergic neurodegeneration due to neuroleptic-induced excitotoxicity and oxidative stress is one of the etiopathological mechanisms in the pathophysiology of tardive dyskinesia and GABA agonists are shown to be effective in ameliorating the symptoms of tardive dyskinesia. The beneficial effect of Ashwagandha root extract might be due to its GABA mimetic activity. Ashwagandha, its constituents and the metabolites of its constituents promote the growth of nerves after taking it for 7 days.

An intriguing study demonstrated that chronic oral administration of withanoside IV attenuated the axonal, dendritic and synaptic losses and memory deficits induced by amyloid peptide A $\beta$ (25–35) in mice. After oral administration in mice, withanoside IV was metabolized into sominone, which induced marked recovery in neurites and synapses and also enhanced axonal and dendritic outgrowth and synaptogenesis. These effects were maintained for at least 7 days after discontinuing withanoside IV administration. These data suggest that withanoside IV, and its metabolite, sominone, may have clinical usefulness as antidementia drugs.

Another team found that the methanol extract of Ashwagandha (5 mg/ml) significantly increased the percentage of cells with neurites in human neuroblastoma SK-N-SH cells. The effect of the extract was dose- and time-dependent. mRNA levels of the dendritic markers MAP2 and PSD-95 by RT-PCR were found to be markedly increased by treatment with the extract. Immunocytochemistry demonstrated the specific expression of MAP2 in neurites extended by the extract. These results suggest that the methanol extract of Ashwagandha promotes the formation of dendrites.

#### Anxiolytic effect

Ashwagandha induced a calming anxiolytic effect that was comparable to the drug Lorazepam in all three standard Anxiety tests: the elevated plus-maze, social interaction and the feeding latency in an unfamiliar environment. Further, both Ashwagandha and Lorazepam, reduced rat brain levels of tribulin, an endocoid marker of clinical anxiety, when the levels were increased following administration of the anxiogenic agent, pentylenetetrazole.

Ashwagandha also exhibited an antidepressant effect, comparable with that induced by imipramine, in two standard tests, the forced swim-induced 'behavioral despair' and 'learned helplessness' tests. The investigations support the use of Ashwagandha as a mood stabilizer in clinical conditions of anxiety and depression.

#### Effect on Energy levels and Mitochondrial Health

The effect of Ashwagandha on glycosaminoglycan synthesis in the granulation tissue of carrageenin-induced air pouch granuloma was studied. Ashwagandha is shown to exert significant inhibitory effect on incorporation of ribosome -35S into the granulation tissue. The uncoupling effect on oxidative phosphorylation (ADP/O ratio reduction) was also observed in the mitochondria of granulation tissue. Further, Mg<sup>2+</sup> dependent ATPase activity was found to be influenced by Ashwagandha. Ashwagandha also reduced the succinate dehydrogenase enzyme activity in the mitochondria of granulation tissue.

#### Anti-inflammatory effect due to Withaferin

Withaferin A and 3-b-hydroxy-2,3-dihydrowithanolide F isolated from *Withania somnifera* show promising antibacterial, antitumoral, immunomodulating and anti-inflammatory properties.

#### Anti-arthritic effect

Ashwagandha is an analgesic that soothes nervous system from pain response. The powerful anti-arthritic properties of Ashwagandha are now widely accepted and documented; it is furthermore found to be effective as antipyretic as well as analgesic also.

Ashwagandha (1000 mg/kg/oral) produced significant analgesic activity for a rat experiencing heat analgesia induced by hot plate method. The peak analgesic effect of Ashwagandha was recorded as 78.03 percent at 2nd hour of administration. The involvement of pain mediators; prostaglandin and 5-hydroxytryptamine in analgesic activity of Ashwagandha was studied by pretreatment with paracetamol (100 mg/kg, ip) and cyproheptadine (10 mg/kg, ip). The analgesic activity of Ashwagandha was potentiated significantly by cyproheptadine, however, paracetamol failed to exhibit any significant change in its activity, suggesting the involvement of serotonin, but not prostaglandins in the analgesic activity of Ashwagandha.

#### DISCUSSION AND CONCLUSION

The available scientific data support the conclusion that Ashwagandha is a real potent regenerative tonic (Rasayana of Ayurveda), due to its multiple pharmacological actions like anti-stress, neuroprotective, antitumor, anti-arthritic, analgesic and anti-inflammatory etc. It is useful for different types of diseases like Parkinson, dementia, memory loss, stress induced diseases, malignoma and others.

Ashwagandha is used as a household remedy by Indians, who consider it as the best tonic for old people and children, and as aphrodisiac by young people. It is one of the best nervine tonics of Ayurveda, the most ancient system of Medical Sciences. Our clinical experience showed that besides the enumerated neurological conditions, brain strokes causing paralysis and neuronal deficit also improve in the long term treatment with Ashwagandha. We are also using it in all forms of cancer including prostate and lung cancers, especially in last stages, giving the patients lot of health benefits. We have some cases of lung cancer who have refused modern therapy and recovered clinically and radiologically with our therapy of Ashwagandha. In a recent seminar on essential drug concept, it was projected as one of the six essential medicinal herbs.

Thus, the above findings clearly indicate that the traditional use of Ashwagandha has a logical and scientific basis. Large scale clinical studies are needed to prove the clinical efficacy of this herb, especially in stress related diseases, neuronal disorders and cancers.

#### REFERENCES

1. Abbas SS, Singh N. *Anti-stress Agents (Herbs) of Indian Origin - Herbal Drugs, A twenty first century perspective*. Delhi: Institute of Nuclear Medicine and Allied Sciences, Defence Research and Development Organization (DRDO), Govt. of India, 2006; 578–591. [Google Scholar]
2. Abbas SS, Singh V, Bhalla M, Singh N. Clinical Study of Organic Ashwagandha in cases of Parkinsonism, Neuropathy, Paralysis and Uterine Tumours (Fibroids and other tumours) including Cutaneous Endodermal Carcinoma; Proc., National Seminar on "Eco-friendly Herbs of Ayurveda in Healthcare of Mankind: A Strategy for Scientific Evaluation and Uniform Standardization" – Lucknow, 2004; 81. [Google Scholar]
3. Abbas SS, Bhalla M, Singh N. *Proc workshop on essential medicines, adverse drug reactions and therapeutic drug monitoring*. Lucknow: Scientific Convention Centre. A clinical study of Organic Ashwagandha in some cases of uterine tumors (fibroids) and dermatofibrosarcoma, 2005; 143–144. [Google Scholar]
4. Abdel-Magied EM, Abdel-Rahman HA, Harraz FM. The effect of aqueous extracts of *Cynomorium coccineum* and *Withania somnifera* on testicular development in immature Wistar rats. *J Ethnopharmacol*, 2001; 75: 1–4. [PubMed] [Google Scholar]
5. Archana R, Namasivayam A. Antistressor effect of *Withania somnifera*. *J Ethnopharmacol*, 1999; 64: 91–93. [PubMed] [Google Scholar]
6. Atta-ur-Rahman, Samina-Abbas, Dur-e-Shahwar, Jamal SA, Choudhary MI, Abbas S. New withanolides from *Withania* spp. *Journal of Natural Products*, 1991; 56: 1000–1006. [Google Scholar]
7. Begum VH, Sadique J. Effect Of *Withania-*

- Somnifera On Glycosaminoglycan Synthesis in Carrageenin-Induced Air Pouch Granuloma. *Biochemical Medicine and Metabolic Biology*, 1987; 38(3): 272–277. [PubMed] [Google Scholar]
8. Bhandari CR. *Ashwagandha (Withania somnifera) "Vanaushadhi Chandrodya" (An Encyclopedia of Indian Herbs)* Vol. 1. Varanasi, India: CS Series of Varanasi Vidyavilas Press, 1970; 96–97. [Google Scholar]
  9. Bhattacharya SK, Goel RK, Kaur R, Ghosal S. Anti-stress activity of Sитоindosides VII and VIII. New Acylsterylglucosides from *Withania somnifera*. *Phytother Res.*, 1987; 1: 32–37. [Google Scholar]
  10. Bhattacharya SK, Kumar A, Ghosal S. Effects of glycowithanolides from *Withania somnifera* on animal model of Alzheimer's disease and perturbed central cholinergic markers of cognition in rats. *Phytother Res.*, 1995; 9: 110–113. [Google Scholar]
  11. Budhiraja RD, Sudhir S. Review of biological activity of withanolides. *JSIR*, 1987; 46: 488–491. [Google Scholar]
  12. Changhadi Govardhan Sharma, author. *Ashwagandharishta - Rastantra Sar Evam Sidhyaprayog Sangrah - Krishna-Gopal Ayurveda Bhawan (Dharmarth Trust)* Nagpur, 1938; 743–744. [Google Scholar]
  13. Charak Samhita 6000BC, author. *Charaka translation into English: Translator: Shree Gulabkunverba Ayurvedic Society*. Jamnagar, India, 1949. [Google Scholar]
  14. Dekosky S, Scheff SW. Synapse loss in frontal cortex biopsies in Alzheimer's disease: correlation with cognitive severity. *Ann Neurol*, 1990; 27: 457–464. [PubMed] [Google Scholar]
  15. Dickon TC, Vickers JC. The morphological phenotype of  $\beta$ -amyloid plaques and associated neuritic changes in Alzheimer's disease. *Neuroscience*, 2001; 105: 99–107. [PubMed] [Google Scholar]
  16. Dixit KS, Agarwal AK, Seth PK, Singh N. *World Congress on Biotech Dev Med Subs Plants & Marine Origin*. Lucknow (India): King George Medical College. Effect of *Withania somnifera*, *Panax ginseng* and *Cannabis indica* and radio ligand binding with neurohumoral in the CNS, 1995; 141. [Google Scholar]
  17. Ghosal S, Srivastava RS, Bhattacharya SK, Upadhyay SN, Jaiswal AK, Chattopadhyay U. Immunomodulatory and CNS effects of sitoindosides IX and X, two new glycowithanolides form *Withania somnifera*. *Phytother Res.*, 1989; 2: 201–206. [Google Scholar]
  18. Gunne LM, Andren PE. An animal model for coexisting tardive dyskinesia and tardive parkinsonism: A glutamate hypothesis for tardive dyskinesia. *Clin Neuropharmacol*, 1993; 16: 90–95. [PubMed] [Google Scholar]
  19. Kalsi R, Singh N, Gupta GP. *Physiology of Human Performance*. Delhi: Defence Institute of Physiology and Allied Sciences, Defence Research and Development Organization (DRDO), Govt. of India. Effects of stress and anti-stress drugs on succinate dehydrogenase enzyme (SDH) in rat brain (A possible role of SDH in stress adaptation phenomenon), 1987; 114–117. [Google Scholar]
  20. Kritikar KR, Basu BD. *Withania somnifera, Indian medicinal plants*. 2nd Edition. IIIrd. Lalit Mohan Basu, Allahabad, 1935; 1774–1776. [Google Scholar]
  21. Kuboyam A T, Tohda C, Komatsu K. Withanoside IV and its active metabolite, sominone, attenuate A beta(25–35)-induced neurodegeneration. *Eur J Neurosci*, 2006; 23(6): 1417–1426. [PubMed] [Google Scholar]
  22. Kulkarni SK, Sharma A, Verma A, Ticku MK. GABA receptor mediated anticonvulsant action of *Withania somnifera* root extract. *Indian Drugs*, 1993; 30: 305–312. [Google Scholar]
  23. Mazen ES, Pavelescu M, Grigorescu E. *Testing of Analgesic Activity of Dichloromethanic and Methanolic Extract from Withania-Somnifera Roots*. 3–4. Vol. 94. Revista Medico-Chirurgicala Societatii de Medici si Naturalisti din Iasi. Contributions to the Pharmacodynamic Study of roots of *Withania somnifera* Dun Species Of Pakistani origin, 1990; 603–605. [PubMed] [Google Scholar]
  24. Misra B. *Ashwagandha - Bhavprakash Nigantu (Indian Materia Medica)* Varanasi: Chaukhambha Bharti Academy, 2004; 393–394. [Google Scholar]
  25. Mishra LC, Singh BB, Dagenais S. Scientific basis for the therapeutic use of *Withania somnifera*. (*Ashwagandha*): A review. *Alternative Medicine Reviews*, 2000; 5: 334–346. [PubMed] [Google Scholar]
  26. Nagashyana N, Sankarankutty P, Nampoothiri MRV, Moahan P, Mohan Kumar P. Association of L- dopa with recovery following Ayurvedic medication in Parkinson's disease. *J Neurol Sci.*, 2000; 176: 1121–1127. [Google Scholar]
  27. Sharma GS. *Ashwagandharishta - Rastantra Sar Evam Sidhyaprayog Sangrah - Krishna-Gopal Ayurveda Bhawan (Dharmarth Trust)* Nagpur, 1938; 743–744. [Google Scholar]
  28. Sharma PV. *Ashwagandha, Dravyaguna Vijana, Chaukhambha Viashwabharti*. Varanasi, 1999; 763–765. [Google Scholar]
  29. Singh N, Agarwal AK, Lata A, Kohli RP. Evaluation of 'adaptogenic' properties of *Withania somnifera*. *Proc Indian Pharmacological Society*, 1976; 17. [Google Scholar]
  30. Singh N, Agarwal AK, Lata A, Kohli RP. *XIIIth Scientific Seminar on Indian Medicine*. Vol. 4. Varanasi: Institute of Medical Sciences. Experimental evaluation of 'adaptogenic' properties of *Withania somnifera*, 1977. [Google Scholar]
  31. Singh N. A new concept on the possible therapy of

- stress disease with 'Adaptogens' (Anti-stress drugs) of indigenous plant origin. *Curr Med Prac*, 1981; 25: 50–55. [Google Scholar]
32. Singh N, Singh SP, Sinha JN, Shanker K, Kohli RP. *Withania somnifera* (Ashwagandha) A rejuvenator herbal drug which enhances survival during stress (An adaptogen) *Int J Crude Drug Res.*, 1982; 3: 29–35. [Google Scholar]
  33. Singh N, Singh SP, Nath C, Kohli RP, Bhargava KP. 5<sup>th</sup> Sepal Congress of Rheumatology. Bangkok. Anti-stress plants as anti-rheumatic agents, 1984; 37. [Google Scholar]
  34. Singh N. A pharmaco-clinical evaluation of some Ayurvedic crude plant drugs as anti-stress agents and their usefulness in some stress diseases of man. *Ann Nat Acad Ind Med*, 1986; 2(1): 14–26. [Google Scholar]
  35. Singh N, Singh SP, Dixit KS, Saxena RC, Kohli RP. A Placebo Controlled Clinical Trial of *Cyprus rotundus*, *Withania somnifera* and their Combination in cases of Rheumatoid Arthritis. *Proc International Seminar*, 1986a; 2: 18–21. [Google Scholar]
  36. Singh N, Singh SP, Nath R, Singh DR, Gupta ML, Kohli RP, Bhargava KP. Prevention of Urethane-induced lung adenomas by *Withania somnifera* (L.) Dunal in albino mice. *Int J Crude Drug Res.*, 1986b; 24(2): 99–100. [Google Scholar]
  37. Singh N. XXXV<sup>th</sup> Annual Congress on Medicinal Plant Research (Freiburg) Effect of *Withania somnifera* and *Panax ginseng* on dopaminergic receptors in rat brain during stress, 1988; 28. [Google Scholar]
  38. Singh N. Stress disease and herbal medicine. *Inaugural issue of J Biotech Med Plant Res, Lucknow*, 1993; 2–4. [Google Scholar]
  39. Singh N. Effect of Anti-stress agents on receptor population in rat brain. *Inaugural issue of J Biotech Med Plant Res, Lucknow*, 1993b; 14. [Google Scholar]
  40. Singh N, Misra N. Experimental methods tools for assessment of anti-stress activity in medicinal plants. *Journal of Biomedical Research*, 1993; 12(182): 124–127. [Google Scholar]
  41. Singh N. *World Congress on Biotech Dev Med Subs Plants & Marine Origin*. Lucknow (India): King George Medical College; Stress disease and herbal medicine, 1995a; 48. [Google Scholar]
  42. Singh N. *Proc Vth WCCN*. China: Anti-stress (Ayurvedic Plants) *Ocimum sanctum* (Tulsi) and *Withania somnifera* (Ashwagandha) in prevention and treatment of cardiovascular disorders, 1995b; 10. [Google Scholar]
  43. Singh N, Singh V, Abbas SS. *Proc 2<sup>nd</sup> World Cong Biotech Dev Herbal Med*. Lucknow (India); Role of Adaptogens / Antistress agents of plant origin in health care & stress diseases of man, 2003; 33. [Google Scholar]
  44. Singh N. *Dilemma of Ayurvedic Herbal Medicines / Food Supplements Proc workshop on Essential Medicines, Adverse Drug Reactions and Therapeutic Drug Monitoring*. Lucknow: Dept. of Pharmacology & Therapeutics, KGMU, Scientific Convention Centre, 2005; 35–41. [Google Scholar]
  45. Singh N. *Ayurvedo-Herbal Medicines - The Need of the Time - Herbal Drugs, A twenty first century perspective*. Delhi: Institute of Nuclear Medicine and Allied Sciences, Defence Research and Development Organization (DRDO), Govt. of India, 2006; 535–547. [Google Scholar]
  46. Singh N. *Herbs - The Life of Man, Need Pharmacoclinical Studies for their Scientific Validation - Relevance of Modern Methods of Pharmacological Studies to Traditional Medicine*. Lucknow: Department of Pharmacology & Therapeutics, C.S.M. Medical University, Scientific Convention Centre, 2008; 37–43. [Google Scholar]
  47. Singh N, Gilca M. *Herbal Medicine - Science embraces tradition - a new insight into the ancient Ayurveda*. Germany: Lambert Academic Publishing, 2010; 51–67. [Google Scholar]
  48. Singh N, Hoette Y, Miller R. *Tulsi 'The Mother Medicine of Nature'*. 2nd Edition. Lucknow: International Institute of Herbal Medicine, 2010; 28–32. [Google Scholar]
  49. Singh RH, Udupa KN. *Clinical and experimental studies on rasayana drugs and rasayana therapy*. New Delhi: Special Research Monograph, Central Council for Research in Ayurveda and Siddha (CCRAS), Ministry of Health and Family Welfare, 1993. [Google Scholar]
  50. Singh RS. *Ashwagandha, Vanaushadhi Nidharsika (Ayurvedic Pharmacopia)* UP Sansthan, 1983; 30–31. [Google Scholar]
  51. Singh SP, Singh DR, Gupta ML, Singh N, Kohli RP. An experimental evaluation of anti-tumor activity of *Withania somnifera* (Ashwagandha) and "Geriforte" *XI Annual Conf IPS Ind Jour Pharmacol*, 1979; 11(1): 65. [Google Scholar]
  52. Sumanran VN, Boddul S, Madhuri D. Differential growth inhibitory effects of *Withania somnifera* root on CHO cells. *Phytother Res.*, 2007; 21: 1–4. [Google Scholar]
  53. Twaij HAA, Elisha EE, Khalid RM. Analgesic studies on some Iraqi medicinal plants. *International Journal of Crude Research*, 1989; 27: 109–112. [Google Scholar]